



[Science Education Patient Care & Advocacy](#)  
[Our Society](#) [Our Mission, Vision & Values](#) [EBMT Membership 2023](#) [Organisational Structure 2023](#) [Financial Highlights 2023](#) [EBMT Partners](#)

## Breadcrumb

1. [Home](#)

## Entity Print



Bertram Glass  
LWP Chair  
Germany

## Major achievements

In 2023 the LWP was able to improve its performance in the most important field of CART therapy.

Two projects – TransCART (CART and allo SCT in 3L treatment of LBCL) and CART in r/r LBCL with CNS involvement – were selected by the GoCART Coalition for funding

as high priority projects. The results of the TransCART were presented at the 49th Annual Meeting of the EBMT in April 2023. IPI at the time of cell therapy was established as an important prognostic tool. CART therapy is especially effective in patients with r/r LBCL and limited adverse prognostic factors. In high-risk r/r LBCL is still a mostly unmet medical need as the relapse incidence (RI) of CART is high (> 60%). Due to its high anti-lymphoma activity but high toxicity allogeneic SCT remains a valuable therapeutic modality e.g. after CART failure.

We performed a retrospective RWE study in patients with r/r LBCL and CNS involvement. With an PFS of 30% and OS of 52% at one year, we could demonstrate that CART in this clinical situation is as effective as in patients without CNS involvement. The results compare favourably to those obtained with other clinical modalities obtained in the same clinical situation. The CART-CNS project was an oral presentation at ASH 2023. A manuscript is preparation.

A third CART related project compared the results of CART treatment with brexucabtagene autoleucel (brexu-cel) in r/r mantle cell lymphoma within the clinical phase II study ZUMA-2 with allogeneic SCT in comparable clinical situation treated by allogeneic SCT registered in the EBMT registry. Brexu-cel was superior in terms of OS (81.3% vs 59.2 % @ 1year) and showed a significantly lower NRM (7.1 vs 21,2 % at 1 year) compared to allogeneic SCT. However, there was no significant difference in PFS and ongoing relapses were seen after brexu-cel whilst the incidence of relapse after two years was every low after allogeneic SCT.

To give a benchmark for further clinical studies and retrospective analysis, we analysed the outcome of autologous and allogeneic SCT in DLBCL. It could show continuously better results for autologous SCT over time but decreasing use with the upcoming CART therapy.

Another group of studies focuses on Hodgkin's lymphoma and the embedding of cellular therapy onto complex treatment algorithms with other modern non-cellular therapies. A joint EBMT / CIBMTR analysis showed that prior use of checkpoint inhibitors significantly improves the results of subsequent allogeneic SCT substantially and may establish a new stand of care for advanced r/r HL.

Ongoing projects focus on the role of CPI and Brentuximab vedotin prior and after autologous SCT.

Another retrospective analysis compared the results of haploidentical versus MUD (8/8 HLA match) allogeneic SCT in lymphomas using post-transplant cyclophosphamide as GVHD prophylaxis. In contrast to the results found e.g. in acute leukaemia, MUD was superior to haploidentical SCT in terms of OS and PFS mostly due to lower incidence of severe GVHD and NRM.

In fields in which the success of new agents is missing, analysis of classical allogeneic and autologous SCT remains of major importance for clinical decision-making. One of these fields is treatment of T cell lymphoma. We started a couple of analysis on the role of allogeneic and autologous SCT in T-NHL and other rare diseases. With the largest cohort of patients analysed so far, we could show that allogeneic stem cell transplantation is of high efficacy. With a 3 years PFS and OS of 48.6% and 55.6% and no relapses beyond 3 years allogeneic SCT was confirmed to be standard of care in r/r NK/T cell lymphoma with curative potential.

## Principal research studies

[Long-term survival after autologous or allogeneic transplantation for T-cell lymphoma: a retrospective analysis of the EBMT-Lymphoma Working Party](#)

[Study type](#)

[Retrospective Studies](#)

[Diseases](#)

[Non-Hodgkin's Lymphoma \(NHL\)](#)

[Group](#)

[Lymphoma Working Party \(LWP\)](#)

[Type of Stem Cell Treatment](#)

[Allogeneic](#)

[Autologous](#)

[Principal investigator](#)

[E. Shumilov](#)

[Outcomes of Autologous and Allogeneic Stem Cell Transplantation in Patients with Hepatosplenic T-cell Lymphoma \(HSTCL\)](#)

[Study type](#)

[Retrospective Studies](#)

[Diseases](#)

[Non-Hodgkin's Lymphoma \(NHL\)](#)

[Group](#)

[Lymphoma Working Party \(LWP\)](#)

[Type of Stem Cell Treatment](#)

[Allogeneic](#)

[Autologous](#)

[Principal investigator](#)

[Imke E. Karsten](#)

[Comparison of the outcomes of patients with mantle cell lymphoma after CAR T cell therapy in the ZUMA-2 collective versus allogeneic stem cell transplantation](#)

[Study type](#)

[Retrospective Studies](#)

[Diseases](#)

[Chronic Myeloid Lymphoma \(CML\)](#)

[Group](#)

[Lymphoma Working Party \(LWP\)](#)

[Type of Stem Cell Treatment](#)

[Allogeneic](#)

[CAR T](#)

[Principal investigator](#)

[Prof. Dr. Sascha Dietrich, Dr. Nora Liebers](#)

[Allogeneic Hematopoietic Cell Transplantation and CAR-T therapy for T-cell/histiocyte-rich Large B-cell Lymphoma \(THRLBCL\) A Retrospective Study of the EBMT Lymphoma Working Party](#)

[Study type](#)

[Retrospective Studies](#)

[Diseases](#)

[Non-Hodgkin's Lymphoma \(NHL\)](#)

[Group](#)

[Lymphoma Working Party \(LWP\)](#)

[Type of Stem Cell Treatment](#)

[Allogeneic](#)

[CAR T](#)

[Principal investigator](#)

[Simon Renders, MD](#)

[Trends in Hematopoietic Stem Cell Transplantation for Patients with Diffuse Large B-Cell Lymphoma. A Retrospective Analysis of the Lymphoma WP of the EBMT](#)

[Study type](#)

[Retrospective Studies](#)

[Diseases](#)

[Non-Hodgkin's Lymphoma \(NHL\)](#)

Group

Lymphoma Working Party (LWP)

Type of Stem Cell Treatment

Allogeneic

Autologous

Principal investigator

Norbert Schmitz, Anna Sureda

Efficacy and toxicity of CAR T cell therapy in patients with primary and secondary central nervous system lymphoma

Study type

Retrospective Studies

Diseases

Non-Hodgkin's Lymphoma (NHL)

Group

GoCART Coalition

Lymphoma Working Party (LWP)

Type of Stem Cell Treatment

CAR T

Principal investigator

Anna Ossami Saïdy

Real life analysis BV use as consolidation therapy in HL with high risk post ASCT

Study type

Retrospective Studies

Diseases

Hodgkin's Disease (HD)

Group

Lymphoma Working Party (LWP)

Type of Stem Cell Treatment

Autologous

Principal investigator

Mariana Bastos Oreiro

Post-transplantation cyclophosphamide-based after HLA-matched versus haploidentical donor transplants in patients with Hodgkin lymphoma: a comparative study of the LWP EBMT

Study type

Retrospective Studies

Diseases

Non-Hodgkin's Lymphoma (NHL)

Group

Lymphoma Working Party (LWP)

Type of Stem Cell Treatment

Allogeneic

Allogeneic - Haploidentical

Autologous

Principal investigator

Dr. Juan Montoro

Outcomes of Allogeneic HCT in patients with Hodgkin Lymphoma in the era of Checkpoint Inhibitors: A joint CIBMTR and EBMT analysis.

Study type

Retrospective Studies

Diseases

Hodgkin's Disease (HD)

Group

Lymphoma Working Party (LWP)

Type of Stem Cell Treatment

Allogeneic

Principal investigator

Miguel-Angel Perales

Allogeneic Stem Cell TransplantationIn NK/ T-cell lymphoma

Study type

Retrospective Studies

Diseases

-

Group

Lymphoma Working Party (LWP)

Type of Stem Cell Treatment

Allogeneic

Principal investigator

Norbert Schmitz

## **Key publications**

2023

Allogeneic hematopoietic stem cell transplantation for NK/T-cell lymphoma: an international collaborative analysis

Group

[Lymphoma Working Party \(LWP\)](#)

[Other society](#)

[1st listed author](#)

[Philipp Berning](#)

[Journal](#)

[Leukemia.](#)

[2023](#)

[Graft-versus-host-disease prophylaxis with ATG or PTCY in patients with lymphoproliferative disorders undergoing reduced intensity conditioning regimen HCT from one antigen mismatched unrelated donor](#)

[Group](#)

[Lymphoma Working Party \(LWP\)](#)

[1st listed author](#)

[Annalisa Paviglianiti](#)

[Journal](#)

[Bone Marrow Transplant.](#)

[2023](#)

[Post-transplant cyclophosphamide-based graft-versus-host disease prophylaxis in HLA-matched and haploidentical donor transplants for patients with Hodgkin lymphoma: a comparative study of the LWP EBMT.: GVHD prophylaxis for patients with Hodgkin lymphoma](#)

[Group](#)

[Lymphoma Working Party \(LWP\)](#)

[1st listed author](#)

[J Montoro](#)

[Journal](#)

[Transplant Cell Ther.](#)

[2023](#)

[Organ complications after CD19 CAR T-cell therapy for large B cell lymphoma: a retrospective study from the EBMT transplant complications and lymphoma working party](#)

[Group](#)

[Transplant Complications Working Party \(TCWP\)](#)

[Lymphoma Working Party \(LWP\)](#)

[1st listed author](#)

[Olaf Penack](#)

[Journal](#)

[Front Immunol.](#)

2023

Haploidentical Versus Matched Unrelated Donor Transplants Using Post-Transplantation Cyclophosphamide for Lymphomas

Group


Lymphoma Working Party (LWP)

1st listed author

Alberto Mussetti

Journal

Transplant Cell Ther.



**2021 2022 2023**

**Oral presentations** 1 2 **10**

**Poster presentations** 0 3 **5**

**Educational events** 1 1 **1**

**Major educational events**





[Event](#)

## **49th Annual Meeting of the FBM**



[Event](#)

## **Lymphoma Working Party Educational Course 2023**

Nov 15, 2023 - Nov 17, 2023 / Warsaw, Poland

Discover more

VISIT THE LWP WEBPAGE